

# Mouse Embryo Assay Certificate of Analysis

ManufacturerMicrotechProductMicropipetteBatchnumber2004011Expiry date04/2023Mouse Embryo TestResultSpecifications100≥ 80 %

### Assay system requested by customer:

1mL of culture medium was placed in a tube with the test article for 30-minutes at 37°C and 5 % CO<sub>2</sub>. Post incubation three 12.5 µl drops of the culture medium was extracted from the test article tube and placed in the corresponding wells of a culture dish; 7 one cell mouse embryos were added to each of the three wells and cultured for 96-hours.

### Control assay method and results:

21 one cell (B<sub>6</sub>C<sub>3</sub>F<sub>1</sub> X B<sub>6</sub>D<sub>2</sub>F<sub>1</sub>) embryos were cultured in triplicate micro drops of culture

21/21 (100%) 1-cell to 2-cell within 24 hr 21/21 (100%) 1-cell to expanded blastocyst within 96 hr

For a valid assay, Embryotech<sup>TM</sup> requires at least 70 % of one cell stage control embryos to develop to expanded blastocyst within 96-hours.

## Test assay method and results:

21 one cell  $(B_6C_3F_1 \ X \ B_6D_2F_1)$  embryos were cultured in triplicate micro drops of culture medium that was extracted from the test article:

21/21 (100%) 1-cell to 2-cell within 24 hr 21/21 (100%) 1-cell to expanded blastocyst within 96 hr

#### **Summary of observations:**

All test and control embryos were selected randomly from a common pool of freshly collected embryos and were cultured in the same incubator at 37°C and 5% CO<sub>2</sub>. 100 percent of the control embryos developed to the expanded blastocyst stage within 96-hours. 100 percent of the test embryos cultured in the extracted culture medium developed to the expanded blastocyst stage within 96-hours.

Release date 27/04/2020

Released by Peggy Kreyser

Signature P.P. P. Mayor

Rev01\_00 / 2019-04-15